

Zelevinsky's involution at roots of unity

Bernard LECLERC*, Jean-Yves THIBON† and Eric VASSEROT‡

1 Février 2008

Abstract

We give a combinatorial algorithm for computing Zelevinsky's involution of the set of isomorphism classes of irreducible representations of the affine Hecke algebra $\hat{H}_m(t)$ when t is a primitive n th root of 1. We show that the same map can also be interpreted in terms of aperiodic nilpotent orbits of $\mathbf{Z}/n\mathbf{Z}$ -graded vector spaces.

1 Introduction

In [32], Zelevinsky has introduced an involution of the Grothendieck group of the category of complex smooth representations of finite length of $G = GL(m, F)$, where F is a p -adic field, and conjectured that this involution permutes the classes of irreducible representations ([32], 9.17). In [33], Zelevinsky further conjectured a geometric description of this involution in terms of the graded nilpotent orbits that parametrize the simple G -modules. Both conjectures have been proved by Mœglin and Waldspurger [22] for the category of admissible representations of G generated by their space of I -fixed vectors, where I is an Iwahori subgroup of G . In this case, by a theorem of Bernstein, Borel and Matsumoto, the conjectures can be reformulated in terms of the Hecke algebra $\hat{H}_m(q)$ of G with respect to I . This is an affine Hecke algebra of type A with parameter q equal to the cardinality of the residue field of F . As shown in [22], the problem then becomes that of describing how the isomorphism classes of simple $\hat{H}_m(q)$ -modules are permuted when the action is twisted by a certain involutive automorphism τ of $\hat{H}_m(q)$. In fact the answer does not depend on q , and is the same for any complex parameter t of infinite multiplicative order.

This is no longer the case if one considers the Hecke algebra $\hat{H}_m(t)$ with t a primitive n th root of 1. According to a conjecture of Vignéras [31], the solution of the problem in this case would be relevant for the l -modular representation theory of G , where l is a prime different from p , and n is the multiplicative order of q in the field with l elements.

In this paper, we obtain results similar to those of Mœglin and Waldspurger in the root of unity case. Namely, (i) given a simple $\hat{H}_m(t)$ -module L , we describe in Section 5.2 a simple combinatorial algorithm for computing the isomorphism class $[L^\tau]$ of the simple module obtained by twisting with τ , and (ii) we prove in Section 5.3 that the geometric

*Département de Mathématiques, Université de Caen, BP 5186, 14032 Caen cedex, France.

†Institut Gaspard Monge, Université de Marne-la-Vallée, Champs-sur-Marne, 77454 Marne-la-Vallée cedex 2, France.

‡Département de Mathématiques, Université de Cergy-Pontoise, 2 Av. A. Chauvin, 95302 Cergy-Pontoise cedex, France.

description conjectured by Zelevinsky still holds in this case, this time in terms of the $\mathbf{Z}/n\mathbf{Z}$ -graded nilpotent orbits parametrizing the simple $\widehat{H}_m(t)$ -modules.

However, our methods are different from those of [22] (and thus we obtain new proofs of these results in the non root of unity case). Our main tool is a recent theorem of Ariki [1] which relates the Grothendieck groups of the algebras $\widehat{H}_m(t)$ with the enveloping algebra $U^-(\mathfrak{g})$, where $\mathfrak{g}$ is the Kac-Moody algebra of type $A_{n-1}^{(1)}$ if t has order n , and A_∞ if t has infinite order. Using this result, we show that our problem is equivalent to describing the natural 2-fold symmetry of Kashiwara's crystal graph of $U_v^-(\mathfrak{g})$, induced by the root diagram automorphism exchanging the simple roots α_i and α_{-i} . Then the combinatorial algorithm for $[L^7]$ follows immediately from the explicit description of this graph (Theorem 4.1), and the equivalent formulation in terms of nilpotent orbits is deduced from a geometric construction of the crystal basis conjectured by Lusztig [18] and recently verified by Kashiwara and Saito [13].

We note that Procter [25] and Aubert [2] have established independently that the first conjecture of Zelevinsky holds for the whole category of complex smooth representations of finite length of G . Assuming this hypothesis, Mœglin and Walspurger had already shown in [22] that their geometric and combinatorial descriptions of Zelevinsky's duality were also valid for this larger class of representations.

The paper is structured as follows. In Section 2 we recall the definition of Zelevinsky's involution, and the parametrization of simple $\widehat{H}_m(t)$ -modules by multisegments. Then we can formulate our problem in a more precise way. In Section 3 we review the Hall-Ringel algebra $\mathcal{H}$ associated with a quiver and the canonical basis of its subalgebra $\mathcal{C}$. In Section 4, using the isomorphism $\mathcal{C} \cong U_v^-(\mathfrak{g})$, we compute the action of the Chevalley generators of $U_v^-(\mathfrak{g})$ on the PBW-basis of $\mathcal{H}$. This allows us to describe the crystal graph of $U_v^-(\mathfrak{g})$ in Lusztig's parametrization by nilpotent orbits, and to derive the combinatorial description of τ (5.2). Finally, in 5.3 we briefly recall the results of Lusztig, Kashiwara and Saito on the construction of the crystal basis in terms of irreducible components of a certain Lagrangian variety, and we deduce from them the geometric description of τ .

2 Affine Hecke algebras

2.1 Let t be a nonzero complex number. The affine Hecke algebra $\widehat{H}_m(t)$ associated to $GL(m)$ is the associative $\mathbf{C}$ -algebra with invertible generators $T_1, \dots, T_{m-1}$ and $y_1, \dots, y_m$ subject to the relations

$$\begin{aligned} T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, & 1 \leq i \leq m-2, \\ T_i T_j &= T_j T_i, & |i-j| > 1, \\ (T_i - t)(T_i + 1) &= 0, & 1 \leq i \leq m-1, \\ y_i y_j &= y_j y_i, & 1 \leq i, j \leq m, \\ y_j T_i &= T_i y_j, & \text{for } j \neq i, i+1, \\ T_i y_i T_i &= t y_{i+1}, & 1 \leq i \leq m-1. \end{aligned}$$

The algebra $\widehat{H}_m(t)$ admits involutive automorphisms $\tau, \flat, \sharp$, given on the generators by

$$\begin{aligned} T_i^\tau &= -t T_{m-i}^{-1}, & y_j^\tau &= y_{m+1-j}, \\ T_i^\flat &= T_{m-i}, & y_j^\flat &= y_{m+1-j}^{-1}, \\ T_i^\sharp &= -t T_i^{-1}, & y_j^\sharp &= y_j^{-1}. \end{aligned}$$

One checks that τ and $\flat$ commute and that $x^\sharp = (x^\flat)^\tau$. The involution $\sharp$ was defined by Iwahori and Matsumoto in [8], p. 280, and τ is usually called Zelevinsky's involution (see [22], p. 146), because of its relation with the involution on representations of $\mathfrak{p}$ -adic $GL(m)$ introduced by Zelevinsky in [32, 33].

2.2 For $\mu = (\mu_1, \dots, \mu_r)$ a composition of m , set $D(\mu) = \{\mu_1 + \dots + \mu_k, 1 \leq k \leq r-1\} = \{d_1, \dots, d_{r-1}\}$ and denote by $\hat{H}_\mu$ the subalgebra of $\hat{H}_m(t)$ generated by $y_1, \dots, y_m$ and $\{T_i \mid i \notin D(\mu)\}$. For $a \in \mathbf{Z}^r$, let $\mathbf{C}_{\mu,a}$ be the 1-dimensional representation of $\hat{H}_\mu$ defined by $T_i \mapsto t$, $i \notin D(\mu)$, and $y_{d_{i-1}+1} \mapsto t^{a_i}$, $i = 1, \dots, r$, where $d_0 = 0$. We denote by $M_{\mu,a}$ the induced $\hat{H}_m(t)$ -module

$$M_{\mu,a} = \hat{H}_m(t) \otimes_{\hat{H}_\mu} \mathbf{C}_{\mu,a}.$$

Let $R(\hat{H}_m(t))$ be the complexified Grothendieck group of the category of finite dimensional $\hat{H}_m(t)$ -modules, and let $R_m(t)$ be the linear span of the classes of the composition factors of the various $M_{\mu,a}$ in $R(\hat{H}_m(t))$. We set

$$R(t) = \bigoplus_{m \geq 0} R_m(t),$$

where we have put for convenience $R_0(t) = \mathbf{C}$. When (ν, b) is a permutation of (μ, a) , *i.e.* $\nu_i = \mu_{\sigma(i)}$ and $b_i = a_{\sigma(i)}$ for some $\sigma \in \mathfrak{S}_r$, the induced modules $M_{\mu,a}$ and $M_{\nu,b}$ are in general non isomorphic, but their classes in $R(t)$ are equal [32] (see [30], p. 455). In the sequel, we shall always work in $R(t)$ and therefore shall only consider pairs (μ, a) up to permutation. In fact, such unordered pairs (μ, a) are naturally identified with certain graded nilpotent orbits O (see below Section 3), which gives a canonical labelling of the classes of the induced modules by these orbits : $[M_{\mu,a}] = [M_O]$. There is a partial order on orbits given by $O \trianglelefteq P$ if $O \subset \overline{P}$.

When t is not a root of unity, it is known [32] that for all O , there is a unique simple module whose class occurs in the expansion of $[M_O]$ but does not occur in any $[M_P]$ with $O \triangleleft P$. Let L_O denote this simple module. Then the L_O are pairwise non isomorphic, and one has in $R(t)$

$$[M_O] = [L_O] + \sum_{O \triangleleft P} K_{O,P} [L_P]$$

for some positive integers $K_{O,P}$. (The $K_{O,P}$ are in fact values at 1 of Kazhdan-Lusztig polynomials of type A , as conjectured by Zelevinsky [33, 34] and proved by Ginzburg [5], Theorem 8.6.23.)

If t is a primitive n th root of unity, we may consider that in a pair (μ, a) , a belongs to $(\mathbf{Z}/n\mathbf{Z})^r$. Let us say that (μ, a) is aperiodic if for each $\ell \in \mathbf{N}^*$, the set $\{a_i \in \mathbf{Z}/n\mathbf{Z} \mid \mu_i = \ell\}$ has at most $n-1$ elements. These are the labels associated to the so-called aperiodic graded nilpotent orbits (see [19], 15.3). It follows from Ariki's theorem [1] that for all aperiodic O , there is a unique simple module L_O whose class occurs in the expansion of $[M_O]$ but does not occur in any $[M_P]$ with $O \triangleleft P$. Moreover, $\{[L_O] \mid O \text{ aperiodic}\}$ is a basis of $R(t)$. Finally, the multiplicities of the $[L_P]$ in the $[M_O]$ are given by certain Kazhdan-Lusztig polynomials of affine type $\tilde{A}$. (For the general type a similar result was previously announced by Grojnowski [7]).

2.3 When t is not a root of unity, the pairs (μ, a) can be identified with Zelevinsky's multisegments [32]. We recall that a segment is an interval $[i, j]$ in $\mathbf{Z}$, and that a multisegment is a formal finite unordered sum $\mathbf{m} = \sum_{i \leq j} m_{ij} [i, j]$. (Here m_{ij} stands for the




Figure 1: The diagrams of $(\mu, a) = ((2, 2, 3, 1, 1, 2, 2, 1), (2, 2, 0, 0, 0, -1, -1, -1))$ and its reduction modulo 2

multiplicity of the segment $[i, j]$ in $\mathbf{m}$.) The multisegment corresponding to (μ, a) is then

$$\mathbf{m} = \sum_{i=1}^r [a_i, a_i + \mu_i - 1].$$

A multisegment can be conveniently regarded as a coloured multipartition, *i.e.* as a sequence $(\lambda^{(i)})_{i \in \mathbf{Z}}$ of partitions, such that the parts of $\lambda^{(i)}$ are the lengths of the segments $[i, j]$, and each cell of the k th column of the Young diagram of $\lambda^{(i)}$ contains the integer $i + k - 1$.

In the case where t is a primitive n th root of unity, we shall also identify the labels (μ, a) with multisegments, this time over $\mathbf{Z}/n\mathbf{Z}$, where we regard $\mathbf{Z}/n\mathbf{Z}$ as a set of n points on the unit circle. For $\ell \in \mathbf{N}^*$ and $i \in \mathbf{Z}/n\mathbf{Z}$ we define the segment of length ℓ and origin i on this circle (or more appropriately the loop) as:

$$[i; \ell] := [i, i + 1, \dots, i + \ell - 1].$$

Sometimes we shall also need the dual notation

$$(\ell; i] := [i - \ell + 1, i - \ell + 2, \dots, i].$$

A multisegment over $\mathbf{Z}/n\mathbf{Z}$ is now a formal finite sum

$$\mathbf{m} = \sum_{i \in \mathbf{Z}/n\mathbf{Z}, \ell \in \mathbf{N}^*} m_{[i; \ell]} [i; \ell] = \sum_{j \in \mathbf{Z}/n\mathbf{Z}, k \in \mathbf{N}^*} m_{(k; j]} (k; j].$$

The cyclic multisegment corresponding to the label (μ, a) is $\mathbf{m} = \sum_i [a_i; \mu_i]$. These cyclic multisegments may also be represented by n -tuples of partitions $(\lambda^{(i)})_{i \in \mathbf{Z}/n\mathbf{Z}}$, with cells labelled by integers modulo n . A label (μ, a) is aperiodic if for each ℓ , there is at least one component $\lambda^{(i)}$ which does not contain the part ℓ (this is, up to conjugation of partitions, the labelling used by Ringel in [29] for a basis of the composition algebra of the cyclic quiver). These definitions are illustrated in Figure 1.

Let $I = \mathbf{Z}$ (*resp.* $\mathbf{Z}/n\mathbf{Z}$). The set of all multisegments over I will be denoted by $\mathcal{M}(I)$, or simply $\mathcal{M}$ if no confusion can arise. Thus we have a parametrization of the simple $\hat{H}_m(t)$ -modules $L_{\mathbf{m}}$ by $\mathbf{m} \in \mathcal{M}(I)$ (with $\mathbf{m}$ aperiodic for $I = \mathbf{Z}/n\mathbf{Z}$). We introduce a $\mathbf{N}^{(I)}$ -grading of $R(t)$ by putting $\deg [L_{\mathbf{m}}] = (d_i)_{i \in I}$, where d_i is the number of cells of the diagram of $\mathbf{m}$ which contain i .

2.4 The three involutions τ , $\flat$ and $\sharp$ induce involutions on the set of simple $\hat{H}_m(t)$ -modules $L_{\mathbf{m}}$, and hence on the set of multisegments $\mathbf{m}$ that parametrize them. It is easy to see that $L_{\mathbf{m}}^{\flat} = L_{\mathbf{m}^{\flat}}$ where if $\mathbf{m} = \sum_{i \in I, \ell \in \mathbf{N}^*} m_{[i; \ell]} [i; \ell]$, then $\mathbf{m}^{\flat} = \sum_{i \in I, \ell \in \mathbf{N}^*} m_{[i; \ell]} (\ell; -i]$.

Therefore, it is equivalent to describe either τ or $\sharp$. For generic t , a geometric description of τ in terms of graded nilpotent orbits has been conjectured by Zelevinsky [33]. This conjecture was proved by Mœglin and Waldspurger [22], who also gave a combinatorial algorithm for computing $\mathbf{m} \mapsto \mathbf{m}^\tau$. An explicit formula for $\mathbf{m}^\tau$ has also been found by Knight and Zelevinsky [15].

The finite Hecke algebra $H_m(t)$ is the quotient of $\widehat{H}_m(t)$ by the relation $y_1 = 1$. Hence, $\sharp$ induces an involution on simple $H_m(t)$ -modules. When t is not a root of unity, these modules are parametrized by partitions of m , and $\sharp$ is just the conjugation of partitions. When t is an n th root of unity, the simple $H_m(t)$ -modules are labelled by n -regular partitions, *i.e.* partitions with no part repeated more than $n - 1$ times, and it was recently proved that $\sharp$ coincides with a bijection previously defined by Mullineux [23]. One proof follows from [16] p. 229 and Ariki's theorem, and a second proof was obtained by Brundan [4] by “quantizing” the results of Kleshchev on modular representations of symmetric groups.

The aim of this paper is to describe the maps τ and $\sharp$ in the affine case when t is a primitive n th root of unity. Proposition 5.1 below can be regarded as a common generalization of Mullineux's bijection as described by Kleshchev [14], and of the algorithm of Mœglin and Waldspurger.

3 Graded nilpotent orbits and Hall-Ringel algebras

3.1 Let Γ denote the quiver of type A_{n-1} (*resp.* A_∞ or $A_{n-1}^{(1)}$). The set of vertices of Γ is $I = [1, n - 1]$ (*resp.* $\mathbf{Z}$ or $\mathbf{Z}/n\mathbf{Z}$), and the set of arrows is $\Omega = \{i \rightarrow i - 1 \mid i, i - 1 \in I\}$. A nilpotent representation of Γ over the field $\mathbf{k}$ is a pair (V, x) , where $V = \bigoplus_{i \in I} V_i$ is an I -graded finite-dimensional vector space, and $x = (x_a)_{a \in \Omega}$ is a degree -1 nilpotent linear operator on V . In other words, x is an element of $N_{V, \Omega}$, the subset of nilpotent elements of

$$E_{V, \Omega} = \bigoplus_{i \rightarrow j \in \Omega} \text{Hom}(V_i, V_j).$$

The vector $\mathbf{d} = \mathbf{dim} V = (\dim V_i)_{i \in I} \in \mathbf{N}^{(I)}$ is the (graded) dimension of the representation. The group $G_V = \prod_{i \in I} GL(V_i)$ acts on $E_{V, \Omega}$ and $N_{V, \Omega}$ by conjugation. Two representations (V, x) and (V, x') are equivalent if x and x' lie in the same orbit. Thus, the set of isomorphism classes of nilpotent Γ -representations is naturally identified with $\mathcal{O} = \sqcup_{\mathbf{d} \in \mathbf{N}^{(I)}} \mathcal{O}_{\mathbf{d}}$, where $\mathcal{O}_{\mathbf{d}}$ is the set of G_V -orbits in $N_{V, \Omega}$ for any graded $\mathbf{k}$ -vector space V of dimension $\mathbf{d}$.

The isomorphism classes of nilpotent Γ -representations are parametrized by $\mathcal{M}(I)$ in the following way. For each vertex $i \in I$, let $\mathbf{k}[i]$ denote the simple Γ -module for which $V = V_i = \mathbf{k}$ and $x = 0$. Given a positive integer ℓ , there is a unique (up to isomorphism) indecomposable Γ -module $\mathbf{k}[\ell; i]$ of length ℓ with head $\mathbf{k}[i]$, and all indecomposable modules are of this type. Therefore, any nilpotent Γ -module M is isomorphic to

$$\mathbf{k}[\mathbf{m}] := \bigoplus_{i \in I, \ell \in \mathbf{N}^*} \mathbf{k}[\ell; i]^{\oplus m_{(\ell; i)}}$$

for a unique multisegment $\mathbf{m}$. We denote by $\mathcal{O}_{\mathbf{m}}$ the G_V -orbit of the module $\mathbf{k}[\mathbf{m}]$. For simplicity, we also denote by $\mathcal{O}_i$ the orbit of the simple module $\mathbf{k}[i]$.

3.2 We recall the definition of the twisted Hall algebra associated with the quiver Γ (see [27, 28, 21]). The classification of nilpotent Γ -modules is independent of the ground field. Take $\mathbf{k} = \mathbf{F}_q$, the field with q elements, and let O, P, Q be orbits in $N_{V,\Omega}, N_{W,\Omega}, N_{U,\Omega}$ respectively, where $\dim U = \dim V + \dim W$. Let M be a Γ -module lying on the orbit Q . Then it is was proved by Ringel [27, 29] that the number of submodules of M with type P and cotype O is a polynomial in q with integer coefficients, independent of the choice of M on Q . This is the Hall polynomial $F_{O,P}^Q(q)$.

Define a bilinear form m on $\mathbf{N}^{(I)}$ by

$$m(\mathbf{a}, \mathbf{b}) = \sum_{i \rightarrow j \in \Omega} a_i b_j + \sum_{i \in I} a_i b_i. \quad (1)$$

Then the generic twisted Hall algebra associated with Γ is the $\mathbf{C}(v)$ -algebra $\mathcal{H}$ with basis $\{u_O \mid O \in \mathcal{O}\}$ and multiplication

$$u_O \circ u_P = v^{m(\dim V, \dim W)} \sum_Q F_{O,P}^Q(v^{-2}) u_Q. \quad (2)$$

(In the setting of [21], u_O stands for the product of characteristic functions of the sets of $\mathbf{F}_{p^{2m}}$ -rational points of O for a fixed prime p .)

3.3 Let O be a G_V -orbit in $N_{V,\Omega}$. One sets

$$\langle O \rangle = v^{\dim O} u_O, \quad (3)$$

where $\dim O$ denotes the dimension of the orbit O (not to be confused with the dimension $\dim V$ of the representation of Γ corresponding to a point of O). The basis $\{\langle O \rangle\}$ is called the Poincaré-Birkoff-Witt basis of $\mathcal{H}$. If P is a G_W -orbit in $N_{W,\Omega}$, it follows from (2) that

$$\langle O \rangle \circ \langle P \rangle = \sum_Q v^{\alpha(O,P,Q)} F_{O,P}^Q(v^{-2}) \langle Q \rangle, \quad (4)$$

where

$$\alpha(O, P, Q) = \dim O + \dim P - \dim Q + m(\dim V, \dim W). \quad (5)$$

In the sequel, we shall use another expression for $\alpha(O, P, Q)$. For $\mathbf{a}, \mathbf{b} \in \mathbf{N}^{(I)}$, set

$$r(\mathbf{a}, \mathbf{b}) = - \sum_{i \rightarrow j \in \Omega} a_i b_j + \sum_{i \in I} a_i b_i, \quad (6)$$

and denote by $\varepsilon(O)$ the dimension of the space of endomorphisms of a module M lying in O . Then

$$\alpha(O, P, Q) = -\varepsilon(O) - \varepsilon(P) + \varepsilon(Q) - r(\dim V, \dim W). \quad (7)$$

Indeed,

$$\dim O = \dim G_V - \dim(\text{Stab } M),$$

and

$$\dim(\text{Stab } M) = \dim(\text{Aut } M) = \dim(\text{End } M) = \varepsilon(O).$$

Thus, if we set $\dim V = (d_{i,V})_{i \in I}$ and $\dim W = (d_{i,W})_{i \in I}$, we have

$$\begin{aligned} \alpha(O, P, Q) &= -\varepsilon(O) - \varepsilon(P) + \varepsilon(Q) + \sum_i d_{i,V}^2 + \sum_i d_{i,W}^2 - \sum_i (d_{i,V} + d_{i,W})^2 + m(\dim V, \dim W) \\ &= -\varepsilon(O) - \varepsilon(P) + \varepsilon(Q) - r(\dim V, \dim W). \end{aligned}$$

3.4 Let $\mathcal{C}$ be the twisted composition algebra of Γ , *i.e.* the subalgebra of $\mathcal{H}$ generated by the characteristic functions $u_i = \langle O_i \rangle$ of the orbits of the simple modules $\mathbf{k}[i]$. When Γ is of type A_{n-1} or A_∞ , one has $\mathcal{C} = \mathcal{H}$, but $\mathcal{C}$ is strictly contained in $\mathcal{H}$ for the type $A_{n-1}^{(1)}$. The vector space $\mathcal{H}$ has a basis $\{b_O \mid O \in \mathcal{O}\}$ given by

$$b_O = \sum_{i, O'} v^{-i + \dim O - \dim O'} \dim \mathcal{H}_{O'}^i(IC_O) \langle O' \rangle \quad (8)$$

where $\mathcal{H}_{O'}^i(IC_O)$ is the stalk at a point of O' of the i th intersection cohomology sheaf of the closure $\overline{O}$ of O .

For Γ of type A_∞ , these sheaves have been first considered by Zelevinsky in [33], where he made a conjecture about their connection with representations of $\mathfrak{p}$ -adic $GL(m)$ known as the $\mathfrak{p}$ -adic analogue of the Kazhdan-Lusztig conjecture. In [34], Zelevinsky further proved that the Poincaré polynomials $\sum_i t^i \dim \mathcal{H}_{O'}^i(IC_O)$ are equal to certain Kazhdan-Lusztig polynomials $P_{w(O')w(O)}(t^2)$ of type A .

For Γ of type $A_{n-1}^{(1)}$, Lusztig has shown that the Poincaré polynomials are also Kazhdan-Lusztig polynomials, this time of affine type $\tilde{A}$ ([17], 11). In [19], he introduced the subfamily of aperiodic nilpotent orbits. These are the orbits $O_{\mathbf{m}}$ parametrized by aperiodic multisegments (see 2.3). It is proved in [20], Section 5 that $\{b_O \mid O \text{ aperiodic}\}$ is a basis of $\mathcal{C}$. Note that non aperiodic orbits O' may occur in the right-hand side of (8) for an aperiodic orbit O .

Let $x \mapsto \bar{x}$ be the ring automorphism of $\mathcal{C}$ defined by $\bar{u}_i = u_i$ and $\bar{v} = v^{-1}$. Let $\mathcal{L}$ be the $\mathbf{Z}[v]$ -submodule of $\mathcal{H}$ spanned by the PBW basis $\langle O_{\mathbf{m}} \rangle$. Let $O \in \mathcal{O}$ and suppose that O is aperiodic if Γ is of type $A_{n-1}^{(1)}$. Then b_O is characterized as the unique element of $\mathcal{L} \cap \mathcal{C}$ such that $\overline{b_O} = b_O$ and $b_O \equiv \langle O \rangle \pmod{v\mathcal{L}}$.

4 Affine Lie algebras and quantum affine algebras

4.1 Let $\mathfrak{g}$ be the Kac-Moody Lie algebra associated with the quiver Γ of type A_{n-1} (*resp.* A_∞ or $A_{n-1}^{(1)}$), namely, $\mathfrak{g} = \mathfrak{sl}_n$ (*resp.* $\mathfrak{g} = \mathfrak{sl}_\infty$ or $\widehat{\mathfrak{sl}}_n$). Let $U_v(\mathfrak{g})$ be the corresponding quantized universal enveloping algebra, with generators $e_i, f_i, v^{h_i}, (i \in I)$. The subalgebra generated by $f_i, (i \in I)$ is denoted by $U_v^-(\mathfrak{g})$. This algebra is isomorphic to the twisted composition algebra $\mathcal{C}$ of Γ , the isomorphism being given by $f_i \mapsto \langle O_i \rangle$. It follows that the twisted Hall algebra $\mathcal{H}$ can be regarded as a left $U_v^-(\mathfrak{g})$ -module. The basis $\{b_O\}$ of $\mathcal{C}$ becomes via this isomorphism a basis of $U_v^-(\mathfrak{g})$: this is Lusztig's canonical basis. It is known that it coincides with Kashiwara's lower global basis of $U_v^-(\mathfrak{g})$. By taking $v = 1$ one obtains the canonical basis of $U^-(\mathfrak{g})$, that we shall still denote by $\{b_O\}$. The aim of this section is to describe Kashiwara's crystal graph of $\mathcal{H}$. To do so, we shall first obtain explicit formulas for the action of the Chevalley generators f_i and their adjoint e'_i on $\mathcal{H}$. We believe that these formulas are of independent interest. For example they yield an algorithm similar to that of [16] for computing the canonical basis in its expansion on the PBW-basis.

4.2 We describe the action of the f_i on $\mathcal{H}$ in the PBW-basis. We assume from now on that Γ is of type $A_{n-1}^{(1)}$. (The other two cases are readily deduced from this one, since $U_v(\mathfrak{sl}_n)$ is a natural subalgebra of $U_v(\widehat{\mathfrak{sl}}_n)$, and $U_v(\mathfrak{sl}_\infty)$ is the “limit $n \rightarrow \infty$ of $U_v(\mathfrak{sl}_n)$ ”).

As explained above, the PBW-basis is naturally labelled by multisegments $\mathbf{m}$ over $\mathbf{Z}/n\mathbf{Z}$. For simplicity, we abuse notation and write $\langle \mathbf{m} \rangle := \langle O_{\mathbf{m}} \rangle$ for the corresponding element of the PBW-basis. Given $i \in \mathbf{Z}/n\mathbf{Z}$, $\ell \in \mathbf{N}^*$, and a multisegment $\mathbf{m}$ such that $m_{(\ell-1; i-1]} \neq 0$ if $\ell > 1$, we define a new multisegment

$$\mathbf{m}_{\ell, i}^+ = \begin{cases} \mathbf{m} + (1; i] & \text{if } \ell = 1, \\ \mathbf{m} + (\ell; i] - (\ell-1; i-1] & \text{if } \ell > 1. \end{cases}$$

In case $m_{(\ell-1; i-1]} = 0$, we put $\langle \mathbf{m}_{\ell, i}^+ \rangle = 0$.

Proposition 4.1 *The Chevalley generators of $U_v^-(\widehat{\mathfrak{sl}}_n)$ act on the PBW-basis of $\mathcal{H}$ by*

$$f_i \langle \mathbf{m} \rangle = \sum_{\ell \in \mathbf{N}^*} v^{\sum_{k > \ell} (m_{(k-1; i-1]} - m_{(k; i]})} [m_{(\ell; i]} + 1] \langle \mathbf{m}_{\ell, i}^+ \rangle,$$

where for $a \in \mathbf{Z}$, $[a] = (v^a - v^{-a})/(v - v^{-1})$ is the usual v -integer.

Proof — We have to apply formulas (4) and (7) with $O = O_i$ and $P = O_{\mathbf{m}}$. We see easily that here, $\varepsilon(O) = 1$, and

$$r(\dim V, \dim W) = \sum_{k \in \mathbf{N}^*} m_{[i; k]} - \sum_{k \in \mathbf{N}^*} m_{(k; i-1]}.$$

Also, it is clear that in this case, the orbits Q giving a nonzero contribution to the right-hand side of (4) are those parametrized by the multisegments of the form $\mathbf{m}_{\ell, i}^+$ for some $\ell \in \mathbf{N}^*$. We distinguish two cases depending on whether $\ell = 1$ or $\ell > 1$.

In the first case we have

$$\begin{aligned} \varepsilon(Q) - \varepsilon(P) &= \dim(\text{End}(\mathbf{k}[i])) + \dim(\text{Hom}(\mathbf{k}[i], \mathbf{k}[\mathbf{m}])) + \dim(\text{Hom}(\mathbf{k}[\mathbf{m}], \mathbf{k}[i])) \\ &= 1 + \sum_{k \in \mathbf{N}^*} m_{[i; k]} + \sum_{k \in \mathbf{N}^*} m_{(k; i]}. \end{aligned}$$

To calculate the Hall polynomial $F_{O, P}^Q$, let us count the number of submodules (V, x) of $(V', x') = \mathbf{F}_q[\mathbf{m}_{1, i}^+]$ isomorphic to $\mathbf{F}_q[\mathbf{m}]$. For $j \neq i$ we must have $V_j = V'_j$, and V_i must be a hyperplane in V'_i which (i) contains $x(V'_{i+1})$ (so that V be stable under x') and (ii) does not contain $\text{Ker } x'|_{V'_i}$ (so that (V, x) be isomorphic to $\mathbf{F}_q[\mathbf{m}]$). The number of hyperplanes satisfying (i) is clearly $\llbracket 1 + \sum_{k \in \mathbf{N}^*} m_{(k; i]} \rrbracket$ and the number of those satisfying (i) but not (ii) is $\llbracket \sum_{k \geq 2} m_{(k; i]} \rrbracket$. (Here we have used the notation $\llbracket a \rrbracket = (q^a - 1)/(q - 1)$.) Therefore,

$$F_{O, P}^Q(q) = \llbracket 1 + \sum_{k \in \mathbf{N}^*} m_{(k; i]} \rrbracket - \llbracket \sum_{k \geq 2} m_{(k; i]} \rrbracket = q^{\sum_{k \geq 2} m_{(k; i]}} \llbracket 1 + m_{(1; i]} \rrbracket,$$

and finally, the coefficient of $\langle \mathbf{m}_{1, i}^+ \rangle$ in $f_i \langle \mathbf{m} \rangle$ is equal to

$$\begin{aligned} v^{-1+1+\sum_{k \in \mathbf{N}^*} m_{[i; k]} + \sum_{k \in \mathbf{N}^*} m_{(k; i]} - \sum_{k \in \mathbf{N}^*} m_{[i; k]} + \sum_{k \in \mathbf{N}^*} m_{(k; i-1]} - 2 \sum_{k \geq 2} m_{(k; i]} - m_{(1; i]}} [m_{(1; i]} + 1] \\ = v^{\sum_{k > 1} (m_{(k-1; i-1]} - m_{(k; i]})} [m_{(1; i]} + 1]. \end{aligned}$$

In the case $\ell > 1$, the calculation is similar and one gets

$$\begin{aligned} \varepsilon(Q) - \varepsilon(P) &= 1 + \sum_{k \in \mathbf{N}^*} m_{[i; k]} + \sum_{k \geq \ell} m_{(k; i]} - \sum_{k \leq \ell-1} m_{(k; i-1]}, \\ F_{O, P}^Q(q) &= \llbracket 1 + \sum_{k \geq \ell} m_{(k; i]} \rrbracket - \llbracket \sum_{k \geq \ell+1} m_{(k; i]} \rrbracket = q^{\sum_{k \geq \ell+1} m_{(k; i]}} \llbracket 1 + m_{(\ell; i]} \rrbracket, \end{aligned}$$

so that the coefficient of $\langle \mathbf{m}_{\ell,i}^+ \rangle$ in $f_i \langle \mathbf{m} \rangle$ is equal to

$$\begin{aligned} & v^{-1+1+\sum_{k \in \mathbf{N}^*} m_{[i;k]} + \sum_{k \geq \ell} m_{(k;i]} - \sum_{k \leq \ell-1} m_{(k;i-1]} - \sum_{k \in \mathbf{N}^*} m_{[i;k]} + \sum_{k \in \mathbf{N}^*} m_{(k;i-1]} \\ & \quad \times v^{-2 \sum_{k \geq \ell+1} m_{(k;i]}} v^{-m_{(\ell;i]}} [m_{(\ell;i]} + 1] \\ & = v^{\sum_{k > \ell} (m_{(k-1;i-1]} - m_{(k;i]})} [m_{(\ell;i]} + 1]. \end{aligned}$$

□

Remark The specialization $v = 1$ of Proposition 4.1 appears in [1] Lemma 4.2.

4.3 To define the Kashiwara operators $\tilde{f}_i$ and $\tilde{e}_i$ of the $U_v^-(\widehat{\mathfrak{sl}}_n)$ -module $\mathcal{H}$, one needs to introduce endomorphisms e'_i of the space $\mathcal{H}$ (see [11], 3.4). First one defines a scalar product on $\mathcal{H}$ for which the basis $\{\langle \mathbf{m} \rangle\}$ is orthogonal by

$$(\langle \mathbf{m} \rangle, \langle \mathbf{m}' \rangle) = v^{-2\varepsilon(\mathbf{m})} \frac{(1-v^2)^{\dim \mathbf{k}[\mathbf{m}]}}{|\text{Aut } \mathbf{F}_q[\mathbf{m}]|_{q=v^{-2}}} \delta_{\mathbf{m}, \mathbf{m}'} . \quad (9)$$

Here we have written for short $\varepsilon(\mathbf{m}) = \varepsilon(O_{\mathbf{m}})$. This scalar product is the one introduced by Green [6], multiplied by a normalization factor $(v^{-2} - 1)^{\dim \mathbf{k}[\mathbf{m}]}$ so that it induces Kashiwara's scalar product [11] on $\mathcal{C} \cong U_v^-(\widehat{\mathfrak{sl}}_n)$.

Lemma 4.1 Put $\varphi_m(t) = (1 - t^m)(1 - t^{m-1}) \cdots (1 - t)$. We have

$$|\text{Aut } \mathbf{F}_q[\mathbf{m}]| = |\text{End } \mathbf{F}_q[\mathbf{m}]| \prod_{i,\ell} \frac{|GL(m_{(\ell;i]}, \mathbf{F}_q)|}{|\text{Mat}(m_{(\ell;i]}, \mathbf{F}_q)|} = q^{\varepsilon(\mathbf{m})} \prod_{i,\ell} \varphi_{m_{(\ell;i]}}(q^{-1}).$$

Proof — Let (V, x) be a Γ -module isomorphic to $\mathbf{F}_q[\mathbf{m}]$, and let $W_{\ell,i}$ denote the subspace of V_i consisting of all the generators of an indecomposable summand of (V, x) of type $(\ell; i]$. Thus, $\dim W_{\ell,i} = m_{(\ell;i]}$.

An endomorphism φ of (V, x) is completely determined by its restriction $\varphi|_W$ to $W = \bigoplus_{\ell,i} W_{\ell,i}$. Let U_i be a complement of $\bigoplus_{\ell} W_{\ell,i}$ in V_i , and $p_{\ell,i}$ be the corresponding projection of V_i onto $W_{\ell,i}$. Then, $x \circ \varphi = \varphi \circ x$ implies that $p_{\ell,i} \circ \varphi|_{W_{k,i}} = 0$ for $\ell > k$. On the other hand, $p_{\ell,i} \circ \varphi|_{W_{\ell,i}}$ is not submitted to any condition and may be an arbitrary element of $\text{End } W_{\ell,i} \cong \text{Mat}(m_{(\ell;i]}, \mathbf{F}_q)$.

Now, φ is an automorphism of (V, x) if and only if $p_W \circ \varphi|_W$ is in $GL(W)$, where $p_W = \sum_{\ell,i} p_{\ell,i}$. Hence φ is an automorphism if and only if $p_{\ell,i} \circ \varphi|_{W_{\ell,i}}$ belongs to $GL(W_{\ell,i}) \cong GL(m_{(\ell;i]}, \mathbf{F}_q)$, and the lemma follows. □

Using Lemma 4.1 and the formula $\dim \mathbf{k}[\mathbf{m}] = \sum_{i,\ell} \ell m_{(\ell;i]}$, we rewrite (9) as

$$(\langle \mathbf{m} \rangle, \langle \mathbf{m}' \rangle) = \prod_{i,\ell} v^{-\binom{m_{(\ell;i]}}{2}} \frac{(1-v^2)^{(\ell-1)m_{(\ell;i]}}}{[m_{(\ell;i]}]!} \delta_{\mathbf{m}, \mathbf{m}'} . \quad (10)$$

Next we define $e'_i \in \text{End}_{\mathbf{C}(v)} \mathcal{H}$ as the adjoint operator of f_i with respect to this scalar product. The following proposition describes the action of e'_i on the PBW-basis. Given $i \in \mathbf{Z}/n\mathbf{Z}$, $\ell \in \mathbf{N}^*$, and a multisegment $\mathbf{m}$ such that $m_{(\ell;i]} \neq 0$, we put

$$\mathbf{m}_{\ell,i}^- = \begin{cases} \mathbf{m} - (1; i] & \text{if } \ell = 1, \\ \mathbf{m} - (\ell; i] + (\ell-1; i-1] & \text{if } \ell > 1. \end{cases}$$

In case $m_{(\ell;i]} = 0$, we put $\langle \mathbf{m}_{\ell,i}^- \rangle = 0$.

Proposition 4.2 *The endomorphisms e'_i act on the PBW-basis of $\mathcal{H}$ by*

$$\begin{aligned} e'_i \langle \mathbf{m} \rangle &= v^{\sum_{k>1} (m_{(k-1;i-1]} - m_{(k;i]})} v^{-m_{(1;i]}+1} \langle \mathbf{m}_{1,i}^- \rangle \\ &+ \sum_{\ell \geq 2} v^{\sum_{k>\ell} (m_{(k-1;i-1]} - m_{(k;i]})} v^{-m_{(\ell;i]}+1} (1 - v^{2(m_{(\ell-1;i-1]}+1)}) \langle \mathbf{m}_{\ell,i}^- \rangle. \end{aligned}$$

Proof — By definition of e'_i and using Proposition 4.1, the coefficient of $\langle \mathbf{m}_{\ell,i}^- \rangle$ in $e'_i \langle \mathbf{m} \rangle$ is equal to

$$\frac{(\langle \mathbf{m} \rangle, f_i \langle \mathbf{m}_{\ell,i}^- \rangle)}{(\langle \mathbf{m}_{\ell,i}^- \rangle, \langle \mathbf{m}_{\ell,i}^- \rangle)} = \frac{(\langle \mathbf{m} \rangle, \langle \mathbf{m} \rangle)}{(\langle \mathbf{m}_{\ell,i}^- \rangle, \langle \mathbf{m}_{\ell,i}^- \rangle)} v^{\sum_{k>\ell} (m_{(k-1;i-1]} - m_{(k;i]})} [m_{(\ell;i]}].$$

Now by (10), if $\ell > 1$

$$\frac{(\langle \mathbf{m} \rangle, \langle \mathbf{m} \rangle)}{(\langle \mathbf{m}_{\ell,i}^- \rangle, \langle \mathbf{m}_{\ell,i}^- \rangle)} = (1 - v^2) v^{-m_{(\ell;i]}+m_{(\ell-1;i-1]}+1} \frac{[m_{(\ell-1;i-1]}+1]}{[m_{(\ell;i]}]},$$

and if $\ell = 1$

$$\frac{(\langle \mathbf{m} \rangle, \langle \mathbf{m} \rangle)}{(\langle \mathbf{m}_{1,i}^- \rangle, \langle \mathbf{m}_{1,i}^- \rangle)} = \frac{v^{-m_{(1;i]}+1}}{[m_{(1;i]}]},$$

which gives the required result. $\square$

Remark Propositions 4.1, 4.2 should be compared to the formulas of [9] for the action of the Chevalley generators of $U_v(\widehat{\mathfrak{sl}}_n)$ on the level l Fock spaces. Actually, our formulas can be regarded as the limit $l \rightarrow \infty$ of those of [9].

4.4 Let $\widehat{B}(\infty)$ be Kashiwara's crystal graph of $\mathcal{H}$ for type $A_{n-1}^{(1)}$ in Lusztig's geometric parametrization. That is, the vertices of this graph are the multisegments $\mathbf{m}$ (or the corresponding orbits $O_{\mathbf{m}}$), and there is an arrow $\mathbf{m} \xrightarrow{i} \mathbf{m}'$ in the graph if and only if $\tilde{f}_i \langle O_{\mathbf{m}} \rangle \equiv \langle O_{\mathbf{m}'} \rangle \pmod{v\mathcal{L}}$, where $\tilde{f}_i$ denotes as usual the Kashiwara operator [11]. In particular, the connected component $B(\infty)$ of this graph containing the empty multisegment gives the crystal graph of $\mathcal{C} \cong U_v^-(\widehat{\mathfrak{sl}}_n)$ parametrized by the set of aperiodic orbits.

There are several descriptions of the crystal graph of $U_v^-(\mathfrak{g})$ available in the literature (see [12], [10], [24]), but the only ones using Lusztig's parametrization are given by Reineke [26] and Kashiwara-Saito [13]. Unfortunately, Reineke's description is restricted to finite dimensional $\mathfrak{g}$, while Kashiwara-Saito's description is in terms of the geometry of orbits, and does not seem to yield a combinatorial algorithm (see Section 5.3). Our description of $\widehat{B}(\infty)$ will be an extension to type $A_{n-1}^{(1)}$ of Reineke's result for type A_{n-1} .

Theorem 4.1 *Let $\mathbf{m}$ be a multisegment over $\mathbf{Z}/n\mathbf{Z}$, and fix $i \in \mathbf{Z}/n\mathbf{Z}$. For $k \in \mathbf{N}$, put $S_{k,i} = \sum_{\ell \geq k} (m_{(\ell;i-1]} - m_{(\ell;i]})$ and let k_0 be minimal such that $S_{k_0,i} = \min_k S_{k,i}$. Then there is an arrow $\mathbf{m} \xrightarrow{i} \mathbf{m}_{k_0,i}^+$ in the crystal graph $\widehat{B}(\infty)$ of $\mathcal{H}$.*

Proof — Our strategy is to reduce the problem to Reineke's description of the crystal graph of $U_v^-(\mathfrak{sl}_n)$. We first note that one can let n tend to infinity in Reineke's work and get the crystal graph of $U_v^-(\mathfrak{sl}_\infty)$.

Let us identify $i \in \mathbf{Z}/n\mathbf{Z}$ with its representative in $\{0, 1, \dots, n-1\}$. Choose n integers $j_0, j_1, \dots, j_{n-1}$ such that $j_r \equiv r \pmod n$ and $j_{i-1} = i-1$, $j_i = i$. For example $\{j_r\} = \{0, 1, \dots, n-1\}$ if $i \neq 0$, and $\{j_r\} = \{-1, 0, 1, \dots, n-2\}$ if $i = 0$. Define an embedding ϕ_i of the set of $\mathbf{Z}/n\mathbf{Z}$ -multisegments in the set of $\mathbf{Z}$ -multisegments by

$$\phi_i \left(\sum_{\ell \in \mathbf{N}^*} \sum_{r \in \mathbf{Z}/n\mathbf{Z}} m_{(\ell; r]}(\ell; r] \right) = \sum_{\ell \in \mathbf{N}^*} \sum_{r \in \mathbf{Z}/n\mathbf{Z}} m_{(\ell; r]}(\ell; j_r].$$

Then define a $\mathbf{C}(v)$ -linear embedding $\Phi_i : \mathcal{H} \longrightarrow U_v^-(\mathfrak{sl}_\infty)$ by

$$\Phi_i(\langle O_{\mathbf{m}} \rangle) = \langle O_{\phi_i(\mathbf{m})} \rangle.$$

Let f_i^∞ denote the Chevalley element in $U_v^-(\mathfrak{sl}_\infty)$ and $e_i^{\prime\infty}$ its adjoint. Proposition 4.1 and 4.2 readily imply

Lemma 4.2 (i) *The subspace $\Phi_i(\mathcal{H})$ of $U_v^-(\mathfrak{sl}_\infty)$ is stable under f_i^∞ and $e_i^{\prime\infty}$.*

(ii) $\Phi_k \circ f_i = f_i^\infty \circ \Phi_i$, $\Phi_i \circ e_i' = e_i^{\prime\infty} \circ \Phi_i$.

Let γ_i (*resp.* γ_i^∞) be the subgraph of the crystal of $\mathcal{H}$ (*resp.* $U_v^-(\mathfrak{sl}_\infty)$) obtained by erasing all arrows labelled by $j \neq i$. Lemma 4.2 shows that

$$\Phi_i \circ \tilde{f}_i = \tilde{f}_i^\infty \circ \Phi_i, \quad \Phi_i \circ \tilde{e}_i = \tilde{e}_i^\infty \circ \Phi_i.$$

Thus, ϕ_i induces an isomorphism of graphs between γ_i and the full subgraph of γ_i^∞ with vertex set $\{\phi_i(\mathbf{m})\}$, where $\mathbf{m}$ runs over all $\mathbf{Z}/n\mathbf{Z}$ -multisegments. The description of this last graph follows from [26] and this finishes the proof of Theorem 4.1. $\square$

Theorem 4.1 is illustrated in Figure 2 in the case $n = 3$.

Remarks 1 It follows easily from Theorem 4.1 that the crystal $\widehat{B}(\infty)$ of $\mathcal{H}$ decomposes into an infinite number of connected components isomorphic to $B(\infty)$, the highest weight vertices of these components being the periodic multisegments, that is, the multisegments $\mathbf{m}$ for which $m_{(\ell; r]} = m_{(\ell; s]}$ for all $r, s \in \mathbf{Z}/n\mathbf{Z}$. This is in agreement with [11], Remarks 3.4.10, 3.5.1.

2 The crystal graphs of the level l integrable representations of $\widehat{\mathfrak{sl}}_n$ as described by [9] embed in a natural way into the crystal graph of $\mathcal{H}$ (*cf.* 4.3 Remark). Each l -tuple of partitions $\boldsymbol{\lambda} = (\lambda^1, \dots, \lambda^l)$ is mapped to the multisegment obtained by taking the formal sum of the rows of the λ^i .

5 Ariki's theorem and Zelevinsky's involution

5.1 Let $K(t)$ denote the graded dual of the $\mathbf{N}^{(I)}$ -graded space $R(t)$. This is made into an associative algebra by taking as product the dual of the comultiplication

$$R_k(t) \rightarrow \bigoplus_{k=k'+k''} R_{k'}(t) \otimes R_{k''}(t)$$

coming from the restriction maps

$$\widehat{H}_k(t) \rightarrow \widehat{H}_{k'}(t) \otimes \widehat{H}_{k''}(t) \quad (k = k' + k'').$$




Figure 2: The crystal graph of $U_v^-(\widehat{\mathfrak{sl}}_3)$ up to degree 3 in its labelling by aperiodic orbits

For any $i \in \mathbf{Z}$ let $\theta_i \in R_1(t)$ be the one-dimensional module such that $\theta_i(y_1) = t^i$. If t is generic (*resp.* $t = e^{2i\pi/n}$) then $\{\theta_i \mid i \in \mathbf{Z}\}$ (*resp.* $\{\theta_i \mid i = 0, 1, \dots, n-1\}$) is a basis of $R_1(t)$. The dual basis vectors are denoted by θ^i . Recall that if M is a simple $\widehat{H}_k(t)$ -module then $c_k := y_1 + y_2 + \dots + y_k$ acts as a scalar, say z_M , on M . Then the i -restriction of the class $[M]$ is defined in [1] as the class $i\text{-res } [M]$ of the $\widehat{H}_{k-1}(t)$ -submodule

$$M_i = \text{Ker}(c_{k-1} - z_M + t^i)^l \subseteq M, \quad l \gg 1,$$

and $i\text{-res}$ is extended to $R(t)$ by linearity. It is straightforward to check that for $M \in R_k(t)$ and $f \in R_{k-1}(t)^*$, one has $(\theta^i \cdot f)(M) = f(i\text{-res}(M))$.

Lemma 5.1 (i) $\sharp$ induces an automorphism of $K(t)$ such that $(\theta^i)^\sharp = \theta^{-i}$.
(ii) τ induces an anti-automorphism of $K(t)$ such that $(\theta^i)^\tau = \theta^i$.

Proof — Claim (i) is obvious since $\sharp$ commutes with the natural embedding

$$i : \widehat{H}_{k'}(t) \otimes \widehat{H}_{k''}(t) \hookrightarrow \widehat{H}_k(t).$$

Claim (ii) follows from the identity $\tau \circ i(a \otimes b) = j(\tau(b) \otimes \tau(a))$ where $a \in \widehat{H}_{k'}(t)$, $b \in \widehat{H}_{k''}(t)$, and j stands for the natural embedding of $\widehat{H}_{k''}(t) \otimes \widehat{H}_{k'}(t)$ into $\widehat{H}_k(t)$. $\square$

5.2 It was proved by Ariki that the map $f_i \mapsto \theta^i$ is an isomorphism of algebras $U^-(\widehat{\mathfrak{sl}}_n) \rightarrow K(t)$ if $t = e^{2i\pi/n}$ (*resp.* $U^-(\mathfrak{sl}_\infty) \rightarrow K(t)$ if t is generic) ([1], Prop. 4.3; see also the Appendix). Recall from 2.2 that $\{[L_O]\}$ is a basis of $R(t)$. Ariki's theorem states that under the previous isomorphism, one has $b_O \mapsto [L_O]^*$, that is, the canonical basis of $U^-(\widehat{\mathfrak{sl}}_n)$ (*resp.* $U^-(\mathfrak{sl}_\infty)$) is mapped to the basis of $K(t)$ dual to $\{[L_O]\}$.




Figure 3: Computing the dual of a multisegment at a 3rd root of 1

Proposition 5.1 *Let $\mathbf{m}$ be a $\mathbf{Z}$ -multisegment (resp. an aperiodic $\mathbf{Z}/n\mathbf{Z}$ -multisegment) and let $t \in \mathbf{C}^*$ be generic (resp. $t = e^{2i\pi/n}$). Let $L_{O_{\mathbf{m}}} = L_{\mathbf{m}}$ be the corresponding simple $\widehat{H}_m(t)$ -module. Then, the twisted module $L_{\mathbf{m}}^{\sharp}$ is isomorphic to $L_{\mathbf{m}^{\sharp}}$, where $\mathbf{m}^{\sharp}$ is the vertex of the crystal graph of $U_v^-(\mathfrak{sl}_{\infty})$ (resp. $U_v^-(\widehat{\mathfrak{sl}}_n)$) obtained from the vertex $\mathbf{m}$ by the 2-fold symmetry $i \longleftrightarrow -i$ of the graph.*

Proof — Let $\sharp$ denote the automorphism of $U^-(\mathfrak{g})$ induced by the 2-fold symmetry $i \longleftrightarrow -i$ of the Dynkin diagram of $\mathfrak{g}$. Repeating the argument of [16] 7.3, 7.4, we see that $\sharp$ preserves the canonical basis $\{b_{\mathbf{m}}\}$ and maps $b_{\mathbf{m}}$ to $b_{\mathbf{m}^{\sharp}}$, where $\mathbf{m}^{\sharp}$ is obtained from $\mathbf{m}$ by the symmetry $i \longleftrightarrow -i$ of the crystal graph. The result then follows from Lemma 5.1 (i) and Ariki's theorem. $\square$

Using Theorem 4.1, it is then easy to describe a simple algorithm for computing $\mathbf{m}^{\sharp}$. Take any path $\emptyset \xrightarrow{i_1} \cdot \xrightarrow{i_2} \cdot \dots \xrightarrow{i_k} \mathbf{m}$ from the empty multisegment to $\mathbf{m}$ in the crystal graph. Then $\emptyset \xrightarrow{-i_1} \cdot \xrightarrow{-i_2} \cdot \dots \xrightarrow{-i_k} \mathbf{m}^{\sharp}$ is a path to $\mathbf{m}^{\sharp}$. This is illustrated in Figure 3 in the case $n = 3$.

Recall that Zelevinsky's involution τ is related to $\sharp$ by $\tau = \sharp \circ \flat$. As explained in 2.4, $\flat$ acts on a multisegment $\mathbf{m}$ by simply changing all segments $(\ell; i]$ into $[-i; \ell)$. Thus Proposition 5.1 gives as well an algorithm for τ .

Remarks 1 Consider the crystal of $U_v^-(\mathfrak{sl}_\infty)$. Among all paths joining a given $\mathbf{m}$ to $\emptyset$, there is a distinguished one obtained by iterating the following procedure. Take the segments $(\ell, i]$ of $\mathbf{m}$ with i minimum, and among them pick up one with ℓ minimum. Then there is always an arrow $\mathbf{m}_{\ell, i}^- \xrightarrow{i} \mathbf{m}$ in the graph. The algorithm for τ obtained by using this particular path is precisely the one described by Moeglin and Waldspurger ([22], Lemme II.3). Note that here, we need a total order on the vertices i of the quiver, and there is no obvious analog of this procedure for the cyclic case.

2 In [3], Berenstein and Zelevinsky have described for $\mathfrak{g} = \mathfrak{sl}_n$ the involutions of the canonical basis of $U^-(\mathfrak{g})$ corresponding to $\sharp$ and τ via Lemma 5.1 and Ariki's theorem. They have related them to the transition map between the two indexations of the basis associated with the two extreme reduced expressions of w_0 ([3], Prop. 3.3). They also noted the coincidence with the multisegment duality coming from the representation theory of p -adic $GL(m)$ (see [3], 3. Remark).

5.3 Let $\mathfrak{g}$ be a symmetrizable Kac-Moody Lie algebra. We briefly review the geometric construction of $B(\infty)$ conjectured by Lusztig ([18], 10.4) and proved by Kashiwara-Saito [13]. Let Γ be the graph associated to the root system of $\mathfrak{g}$. Let I be the set of vertices. Fix an orientation Ω of Γ and set $H = \Omega \cup \overline{\Omega}$, where $\overline{\Omega}$ denotes the orientation opposite to Ω . For any arrow $a \in H$ let $\text{in}(a)$ and $\text{out}(a)$ be the input and output vertices of a . Given a I -graded complex vector space $V = \bigoplus_{i \in I} V_i$, Lusztig has introduced [18, 19]

$$X_V = \bigoplus_{a \in H} \text{Hom}(V_{\text{in}(a)}, V_{\text{out}(a)}),$$

and as above,

$$E_{V, \Omega} = \bigoplus_{a \in \Omega} \text{Hom}(V_{\text{in}(a)}, V_{\text{out}(a)}).$$

The space $E_{V, \Omega}^*$ is naturally identified with $E_{V, \overline{\Omega}}$, and, thus, $X_V = E_{V, \Omega} \oplus E_{V, \overline{\Omega}}$ gets identified with the cotangent bundle of $E_{V, \Omega}$. It is therefore a symplectic manifold, and the natural action of G_V on X_V is Hamiltonian, with moment map $\mu : X_V \rightarrow \mathfrak{g}_V$, where $\mathfrak{g}_V$ is the Lie algebra of G_V . The i -th component of μ is the map

$$\mu_i : X_V \rightarrow \text{End}(V_i), \quad B \mapsto \sum_{\substack{a \in H \\ \text{in}(a)=i}} \epsilon(a) B_{\overline{a}} B_a,$$

where $\epsilon(a) = 1$ if $a \in \Omega$ and -1 if $a \in \overline{\Omega}$. The set $\{B \in X_V \mid \mu(B) = 0\}$ is easily seen to be the union of the conormal bundles C_O of the G_V -orbits $O \in E_{V, \Omega}$.

An element $B \in X_V$ is nilpotent if there is an integer l such that $B_{a_1} \circ B_{a_2} \circ \dots \circ B_{a_l} = 0$ for all sequences $(a_1, a_2, \dots, a_l)$ such that the above expression is meaningful. Define

$$\Lambda_V = \{B \in X_V \mid \mu(B) = 0 \text{ and } B \text{ is nilpotent}\}.$$

It is known [18, 19] that Λ_V is a closed equidimensional G_V -stable subvariety of X_V . For Γ of type A_{n-1} , A_∞ or $A_{n-1}^{(1)}$, an element B of Λ_V can be identified with a pair (x, y) of commuting nilpotent endomorphisms of V of degree -1 and $+1$ respectively. Moreover,

the irreducible components of Λ_V are the closures $\overline{CO}$ of the conormal bundles to the nilpotent G_V -orbits O in $E_{V,\Omega}$ for type A_{n-1} or A_∞ ([19], 14), and to the aperiodic nilpotent G_V -orbits for type $A_{n-1}^{(1)}$ ([19], 15).

To the I -graded space V we associate the negative root $\nu = -\sum_i (\dim V_i) \alpha_i$. Let $B(\infty; \nu)$ be the set of irreducible components of Λ_V . Lusztig has introduced a crystal structure on the set $\sqcup_\nu B(\infty; \nu)$, and conjectured that this graph is isomorphic to the crystal $B(\infty)$ of $U_v^-(\mathfrak{g})$ ([18], 8.9). This was proved by Kashiwara-Saito [13]. More precisely, recall that the vertices of $B(\infty)$ are identified with orbits $O \in \mathcal{O}$ for type A_{n-1} or A_∞ , and aperiodic orbits $O \in \mathcal{O}$ for type $A_{n-1}^{(1)}$. Then the map $\overline{CO} \mapsto O$ is an isomorphism of crystals $\sqcup_\nu B(\infty; \nu) \rightarrow B(\infty)$.

The algebra $U_v(\mathfrak{g})$ admits an anti-automorphism $*$ such that $e_i^* = e_i$, $f_i^* = f_i$, and $(v^h)^* = v^{-h}$, which induces an involution (still denoted by $*$) of $B(\infty)$ [11]. It is shown in [13], Section 5.3 that the involution of $B(\infty; \nu)$ induced by the transpose map $B \mapsto {}^t B$ on Λ_V coincides with $*$. (This was also conjectured by Lusztig [18], 10.4.) More precisely, if $x \in O \subseteq N_{V,\Omega}$ then $*(\overline{CO})$ is the irreducible component containing $({}^t y, {}^t x)$, where $y \in N_{V,\overline{\Omega}}$ is generic such that $[x, y] = 0$. Summarizing the discussion, and taking into account Lemma 5.1 (ii) and Ariki's theorem, we can state:

Proposition 5.2 *Let Γ be of type $A_{n-1}^{(1)}$ or A_∞ . For x in $E_{V,\Omega}$, set*

$$Z(x) = \{y \in E_{V,\overline{\Omega}} \mid x \circ y = y \circ x\}.$$

Assume that x is nilpotent (and aperiodic in type $A_{n-1}^{(1)}$). Then,

- (i) *there is a unique G_V -orbit $O' \subset N_{V,\overline{\Omega}}$ such that $O' \cap Z(x)$ is Zarisky dense and open in $Z(x)$, and this orbit only depends on the orbit O of x ;*
- (ii) *if $O = O_{\mathbf{m}}$ and $y \in O'$, then ${}^t y \in O_{\mathbf{m}^\tau}$, where the multisegment $\mathbf{m}^\tau$ is obtained from $\mathbf{m}$ as explained in 5.2.*

For Γ of type A_∞ , we recover the description of Zelevinsky's involution conjectured in [33] and established in [22].

6 Appendix

In [1] Ariki uses the non-vanishing theorem for affine Hecke algebras at roots of unity announced in [7]. The purpose of this appendix is to indicate a simple proof of this non-vanishing result for type A . S. Ariki has informed one of us (E.V.) that this simple argument was known to him and to G. Lusztig. For the sake of completeness, we include it here.

Fix $m \in \mathbf{N}^*$ and put $t = e^{2i\pi/n}$. We use the same notations as in Sections 3.1 and 5.3. In particular (Γ, Ω) is the cyclic quiver with n vertices and V is a I -graded vector space of dimension $\mathbf{d} \in \mathbf{N}^{(I)}$. Let $S_{\mathbf{d}}$ be the set of sequences $\mathbf{i} = (i_1, i_2, \dots, i_m) \in I^m$ such that $\sharp\{k \mid i_k = i\} = d_i$. Given such a sequence let $F_{\mathbf{i}}$ be the variety of flags of type $\mathbf{i}$ in V (see [19], Section 1). Let $\tilde{F}_{\mathbf{i}}$ be the variety of pairs $(\phi, x) \in F_{\mathbf{i}} \times E_{V,\Omega}$ such that ϕ is x -stable. As usual, we denote by $\mathcal{D}(X)$ the bounded derived category of complexes of $\mathbf{C}$ -sheaves on a complex algebraic variety X . Let $\pi_{\mathbf{i}} : \tilde{F}_{\mathbf{i}} \rightarrow E_{V,\Omega}$ be the second projection and put $\mathcal{L}_{\mathbf{i}} = (\pi_{\mathbf{i}})_!(\mathbf{C}) \in \mathcal{D}(E_{V,\Omega})$. By definition the degree- $\mathbf{d}$ component of Lusztig's canonical basis $\{b_O\}$ is labelled by the G_V -orbits $O \subset N_{V,\Omega}$ such that a shift of the intersection cohomology complex IC_O is a direct factor in $\bigoplus_{\mathbf{i} \in S_{\mathbf{d}}} \mathcal{L}_{\mathbf{i}}$ (use [19], Section 2.4).

Let F be the variety of complete flags in $\mathbf{C}^m$. The cotangent variety T^*F is identified with the set of pairs $(\phi, x) \in F \times \text{End}(\mathbf{C}^m)$ such that ϕ is x -stable. If $s \in GL(m)$ is semi-simple let $G_s \subseteq GL(m)$ be its centralizer, let $N_s \subset \text{End}(\mathbf{C}^m)$ be the set of nilpotent matrices x such that $sxs^{-1} = t^{-1}x$ and let $\tilde{F}_s$ be the set of pairs $(\phi, x) \in T^*F$ such that $x \in N_s$ and the flag ϕ is fixed by s . Let $\pi_s : \tilde{F}_s \rightarrow N_s$ be the second projection and set $\mathcal{L}_s = (\pi_s)_!(\mathbf{C}) \in \mathcal{D}(N_s)$. Given a sequence $\mathbf{d} \in \mathbf{N}^{(I)}$ such that $\sum_i d_i = m$, if s is a diagonal matrix such that t^i has the multiplicity d_i in the spectrum of s then G_s, N_s are identified with $G_V, N_{V,\Omega}$ above. In this case set $\mathcal{L}_{\mathbf{d}} = \mathcal{L}_s$. Then, Ginzburg's geometric construction of the simple modules of the affine Hecke algebra (see [5]) implies that the simple modules of $\hat{H}_m(t)$ are labelled by the G_V -orbits $O \subset N_{V,\Omega}$ such that a shift of the complex IC_O is a direct factor in $\bigoplus_{\mathbf{d}} \mathcal{L}_{\mathbf{d}}$.

In our case the non-vanishing result is precisely that the orbits labelling the degree- m component of the basis $\{b_O\}$ are the same as the orbits labelling the simple $\hat{H}_m(t)$ -modules. It is a consequence of the obvious equality of the complexes $\mathcal{L}_{\mathbf{d}}$ and $\bigoplus_{\mathbf{i} \in S_{\mathbf{d}}} \mathcal{L}_{\mathbf{i}}$ (simply observe that $F_{\mathbf{d}} = \sqcup_{\mathbf{i} \in S_{\mathbf{d}}} F_{\mathbf{i}}$).

Acknowledgements We are grateful to M.-F. Vigneras for posing the question which motivated this work. B.L. and J.-Y.T. would like to thank A. Zelevinsky for his beautiful series of lectures on “Representations of quantum groups and piecewise linear combinatorics” given at the Université de Marne-la-Vallée in 1994, which arose their interest in this field.

E.V. is partially supported by EEC grant no ERB FMRX-CT97-0100.

References

- [1] S. ARIKI, *On the decomposition numbers of the Hecke algebra of $G(n, 1, m)$* , J. Math. Kyoto Univ. **36** (1996), 789–808.
- [2] A.-M. AUBERT, *Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d'un groupe p -adique*, Trans. A.M.S., **347** (1995), 2179–2189; Erratum Trans. A.M.S., **348** (1996), 4687–4690.
- [3] A. BERENSTEIN and A. ZELEVINSKY, *Canonical bases for the quantum group of type A_r and piecewise-linear combinatorics*, Duke Math. J. **82** (1996), 473–502.
- [4] J. BRUNDAN, *Modular branching rules and the Mullineux map for Hecke algebras of type A* , to appear in Proc. London Math. Soc..
- [5] N. CHRISS and V. GINZBURG, *Representation theory and complex geometry*, Birkhauser 1997.
- [6] J.A. GREEN, *Hall algebras, hereditary algebras and quantum groups*, Invent. Math. **120** (1995), 361–377.
- [7] I. GROJNOWSKI, *Representations of affine Hecke algebras (and affine quantum GL_n) at roots of unity*, Internat. Math. Res. Notices **5** (1994), 215–217.
- [8] N. IWAHORI and H. MATSUMOTO, *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups*, Publ. Math. I.H.E.S. **25** (1965), 237–280.
- [9] M. JIMBO, K. MISRA, T. MIWA and M. OKADO, *Combinatorics of representations of $U_q(\widehat{\mathfrak{sl}}(n))$ at $q = 0$* , Commun. Math. Phys. **136** (1991), 543–566.
- [10] S.-J. KANG, M. KASHIWARA and K. MISRA, *Crystal bases of Verma modules for quantum affine algebras*, Compositio Mathematica **92** (1994), 299–325.
- [11] M. KASHIWARA, *On crystal bases of the q -analogue of universal enveloping algebra*, Duke Math. J. **63** (1991), 465–516.
- [12] M. KASHIWARA, *Crystal base and Littlemann's refined Demazure character formula*, Duke Math. J. **71** (1993), 839–858.

- [13] M. KASHIWARA and Y. SAITO, *Geometric construction of crystal bases*, Duke Math. J. **89** (1997), 9–36.
- [14] A.S. KLESHCHEV, *Branching rules for the modular representations of symmetric groups III; some corollaries and a problem of Mullineux*, J. London Math. Soc., **2** (1995).
- [15] H. KNIGHT and A. ZELEVINSKY, *Representations of quivers of type A and the multisegment duality*, Adv. Math. **117** (1996), 273–293.
- [16] A. LASCoux, B. LECLERC and J.-Y. THIBON, *Hecke algebras at roots of unity and crystal bases of quantum affine algebras*, Commun. Math. Phys. **181** (1996), 205–263.
- [17] G. LUSZTIG, *Canonical bases arising from quantized enveloping algebras, I*, J. Amer. Math. Soc. **3** (1990), 447–498.
- [18] G. LUSZTIG, *Canonical bases arising from quantized enveloping algebras, II*, Progress Theor. Phys. Supplement **102** (1990), 175–201.
- [19] G. LUSZTIG, *Quivers, perverse sheaves and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), 365–421.
- [20] G. LUSZTIG, *Affine quivers and canonical bases*, Publ. Math., I.H.E.S. **76** (1992), 111–163.
- [21] G. LUSZTIG, *Canonical bases and Hall algebras*, Lectures at the Summer School on Representation Theory and Algebraic Geometry, Montréal 1997.
- [22] C. MOEGLIN and J.-L. WALDSPURGER, *Sur l’involution de Zelevinski*, J. Reine Angew. Math. **372** (1986), 136–177.
- [23] G. MULLINEUX, *Bijections of p -regular partitions and p -modular irreducibles of the symmetric groups*, J. London Math. Soc., **20** (1979), 60–66.
- [24] T. NAKASHIMA and A. ZELEVINSKY, *Polyhedral realizations of crystal bases for quantized Kac-Moody algebras*, Advances Math. **131** (1997), 253–278.
- [25] K. PROCTER, *The Zelevinsky duality conjecture for GL_n* , Thesis King’s College, London, 1994.
- [26] M. REINEKE, *On the coloured graph structure of Lusztig’s canonical basis*, Math. Ann. **307** (1997), 705–723.
- [27] C. M. RINGEL, *Hall algebras*, in *Topics in Algebra*, Banach Center Publ. **26** (1990), 433–447.
- [28] C. M. RINGEL, *Hall algebras revisited*, in *Quantum deformations of algebras and their representations*, Israel Math. Conf. Proc., Ed. A. Joseph, S. Shnider, **7** (1993), 171–176.
- [29] C. M. RINGEL, *The composition algebra of a cyclic quiver*, Proc. London Math. Soc. **66** (1993), 507–537.
- [30] J. D. ROGAWSKI, *On modules over the Hecke algebra of a p -adic group*, Invent. Math. **79** (1985), 443–465.
- [31] M. F. VIGNERAS, *A propos d’une conjecture de Langlands modulaire*, in *Finite reductive groups: related structures and representations*, Cabanes, Marc (ed.), Birkhauser 1997.
- [32] A. ZELEVINSKY, *Induced representations of reductive p -adic groups II*, Ann. Sci. E.N.S. **13** (1980), 165–210.
- [33] A. ZELEVINSKY, *A p -adic analog of the Kazhdan-Lusztig conjecture*, Funct. Anal. Appl. **15** (1981), 83–92.
- [34] A. ZELEVINSKY, *Two remarks on graded nilpotent classes*, Russian Math. Surveys **40** (1985), 249–250.